

## Meds for Opioid Overdose

To reverse an opioid overdose in the community setting, opioid receptor blockers can be administered by nasal spray (naloxone, nalmefene) or IM injection (naloxone). In the hospital setting naloxone or nalmefene are usually administered parenterally. The FAQ below addresses common questions regarding these opioid receptor blockers. See our *Naloxone Quick Start Guide* for a simplified resource about identifying patients for whom naloxone may be beneficial and getting them started.

Question	Answer/Pertinent Information/Suggested Resources
<b>Who should an opioid receptor blocker be considered for?</b>	<ul style="list-style-type: none"> <li>• Consider for people:               <ul style="list-style-type: none"> <li>○ with a history of opioid intoxication or overdose.<sup>1,9,27</sup></li> <li>○ with a suspected history of substance abuse or nonmedical opioid use (e.g., opioid use disorder).<sup>1,9,27</sup></li> <li>○ on treatment (buprenorphine, etc) for opioid use disorder.<sup>1,9,27</sup></li> <li>○ taking high opioid doses (e.g., 50 mg or more of oral morphine or its equivalent) daily.<sup>18,27</sup> <ul style="list-style-type: none"> <li>▪ Tools to calculate daily morphine equivalents can be found at:                   <ul style="list-style-type: none"> <li>• <a href="https://stacks.cdc.gov/view/cdc/38481">https://stacks.cdc.gov/view/cdc/38481</a>.</li> <li>• <a href="http://www.agencymeddirectors.wa.gov/Calculator/DoseCalculator">http://www.agencymeddirectors.wa.gov/Calculator/DoseCalculator</a>.</li> <li>• <a href="https://www.cdc.gov/opioids/providers/prescribing/pdf/Opioid-Prescribing-Guideline-MobileApp.pdf">https://www.cdc.gov/opioids/providers/prescribing/pdf/Opioid-Prescribing-Guideline-MobileApp.pdf</a> (app also includes helpful opioid prescribing info).</li> </ul> </li> </ul> </li> <li>○ being rotated from one opioid to another (due to risk of incomplete cross-tolerance).<sup>9</sup></li> <li>○ taking an opioid who:<sup>1,9,27</sup> <ul style="list-style-type: none"> <li>▪ smoke or have a respiratory illness (e.g., COPD, sleep apnea, asthma).</li> <li>▪ have kidney, liver, or heart disease; or human immunodeficiency virus (HIV).</li> <li>▪ use alcohol or a benzodiazepine, sedative, or antidepressant.</li> </ul> </li> <li>○ who live in a remote location (consider providing more than one kit or dose due to potential for delays with emergency response personnel in remote areas).<sup>9</sup></li> <li>○ who live with people at risk for accidental opioid overdose (e.g., children).<sup>1</sup></li> <li>○ who request it.<sup>9</sup></li> <li>○ who may be in a position to help some at risk of an opioid overdose.<sup>27</sup></li> </ul> </li> <li>• The FDA is requiring that recommendations for co-prescription of naloxone be added to the prescribing information of opioid pain medicine, as well as medicines used to treat opioid use disorder.<sup>1</sup></li> <li>• In some states, naloxone co-prescription is required for certain high-risk patients.<sup>15</sup></li> </ul>

Question	Answer/Pertinent Information/Suggested Resources
<b>What should you tell patients, caregivers, and families about naloxone and nalmefene?</b>	<ul style="list-style-type: none"><li>• Help patients, caregivers, and families <b>get on board</b> with naloxone and nalmefene by covering these talking points:<ul style="list-style-type: none"><li>○ Anyone on an opioid can be at risk of respiratory depression. Naloxone or nalmefene is like a seatbelt: most people don't need it, but it's there just in case they need it.</li><li>○ Reinforce that naloxone or nalmefene can <b>save a life</b>.</li><li>○ Naloxone or nalmefene are easy to use, have low risk of adverse effects, and are not harmful if the person didn't really need it.</li><li>○ Note that the vast majority of patients are <b>not offended</b> by an offer of naloxone or nalmefene.<sup>12</sup></li></ul></li><li>• Help patients, caregivers, and families <b>get</b> naloxone or nalmefene. Consider keeping an updated list of free naloxone sources in your community, what insurance might cover, and manufacturer programs to cover costs.</li><li>• Help patients, caregivers, and families <b>get trained</b> to administer naloxone.<ul style="list-style-type: none"><li>○ US: <a href="https://prescribeto prevent.org/patient-education/materials/">https://prescribeto prevent.org/patient-education/materials/</a>.</li><li>○ Video from the Canadian Pharmacists Association (Naloxone Made Easy): <a href="https://www.youtube.com/watch?v=0Z-y7CoeDMc">https://www.youtube.com/watch?v=0Z-y7CoeDMc</a>.</li><li>○ Training will cover the essential steps to take if an opioid overdose is suspected: check for signs of opioid overdose (reduced level of consciousness with breathing difficulty or blue/purple lips or nails), try to get person to respond (e.g., shout the person's name, sternal rub), call 911, provide rescue breathing if needed, and give naloxone.<sup>6,17</sup></li><li>○ Patient instruction sheets from the College of Pharmacists of British Columbia are available at <a href="https://www.bcpharmacists.org/naloxone">https://www.bcpharmacists.org/naloxone</a>.</li></ul></li><li>• Explain to family and caregivers <b>what to expect</b> after naloxone or nalmefene administration.<ul style="list-style-type: none"><li>○ Most patients respond to naloxone and return to spontaneous breathing with only mild withdrawal symptoms.<sup>6</sup></li><li>○ Opioid withdrawal is not typically life-threatening in adults.<sup>6,34</sup></li><li>○ <b>Opioid withdrawal symptoms</b> can include sweating, goose bumps, achiness, shivering, gastrointestinal symptoms, tachycardia, irritability, increased blood pressure, fever, runny nose, sneezing, trembling, and yawning.<sup>6,34</sup><ul style="list-style-type: none"><li>▪ Patients may become agitated or confused, or vomit after naloxone or nalmefene is given.<sup>6,34</sup></li></ul></li><li>○ To <b>prevent aspiration</b>, the patient should be positioned on their side after naloxone or nalmefene is given.<sup>6,34</sup><ul style="list-style-type: none"><li>▪ This "recovery position" is illustrated in the <i>Narcan</i>, <i>Kloxxado</i>, and <i>Opvee</i> nasal sprays, <i>Zimhi</i>, and <i>S.O.S Naloxone</i> patient labeling, and in the patient instruction sheets available at <a href="http://www.bcpharmacists.org/naloxone">http://www.bcpharmacists.org/naloxone</a>.</li></ul></li></ul></li></ul>

Question	Answer/Pertinent Information/Suggested Resources
<p><b>When should an opioid receptor blocker be administered?</b></p>	<ul style="list-style-type: none"> <li>• Naloxone or nalmefene should be given if a person has respiratory and/or CNS depression in a situation where opioids may be present.<sup>3,16,25,26,32,34</sup> <ul style="list-style-type: none"> <li>○ Give if the patient is excessively sleepy and cannot be aroused with a loud voice or sternal rub.<sup>3,25,26,32</sup></li> <li>○ Other indications include slow, shallow, or no respirations, or pinpoint pupils in a patient who is difficult to arouse.<sup>3,25,26,32</sup></li> <li>○ Other signs of overdose include blue or purple fingernails or lips, death rattle (gurgling noise in the throat from build-up of saliva and mucus), slow heartbeat, or low blood pressure.<sup>3,6,16</sup></li> </ul> </li> </ul>
<p><b>What is the role of nalmefene nasal spray (<i>Opvee</i>, US only) in opioid overdose?</b></p>	<ul style="list-style-type: none"> <li>• <b>Nalmefene</b> (available by prescription only) is an opioid antagonist that may be used as a naloxone alternative.</li> <li>• It is indicated for the emergency treatment of known or suspected opioid overdose (respiratory and/or CNS depression) in patients <b>12 years and older</b>.<sup>34</sup></li> <li>• Intranasal nalmefene has a similar onset (three to five minutes) and a longer duration of action (six hours) compared to naloxone.<sup>7,24,31</sup></li> <li>• There is no good evidence that nalmefene is more effective than naloxone for opioid overdose. Its longer duration of action theoretically could reduce the risk of relapse of opioid overdose symptoms, but could also prolong opioid withdrawal symptoms.</li> <li>• Check your local state laws and procedures. Some states allow pharmacists to provide nalmefene (as they do with naloxone) by standing orders, protocols, etc.<sup>35,36</sup></li> </ul>
<p><b>Is naloxone available without a prescription?</b></p>	<ul style="list-style-type: none"> <li>• <b>US:</b> Naloxone is available without a prescription.<sup>29</sup> <ul style="list-style-type: none"> <li>○ <b>Narcan nasal spray 4 mg</b> has been FDA-approved for Rx-to-OTC switch. It will be available in pharmacies, grocery stores, gas stations, and other retail and online stores.<sup>33</sup></li> <li>○ To understand specific laws about prescription naloxone in your state (e.g., protocol, standing order), consult your state pharmacy board, or go to one of these websites: <ul style="list-style-type: none"> <li>▪ SafeProject: <a href="https://www.safeproject.us/naloxone-awareness-project/state-rules/">https://www.safeproject.us/naloxone-awareness-project/state-rules/</a>.</li> <li>▪ National Alliance of State Pharmacy Associations: <a href="https://naspa.us/resource/naloxone-access-community-pharmacies/">https://naspa.us/resource/naloxone-access-community-pharmacies/</a>.</li> <li>▪ Prescription Drug Abuse Policy System <a href="https://pdaps.org/datasets/laws-regulating-administration-of-naloxone-1501695139">https://pdaps.org/datasets/laws-regulating-administration-of-naloxone-1501695139</a>.</li> </ul> </li> </ul> </li> <li>• <b>Canada:</b> Naloxone is available without a prescription (unscheduled in Alberta, British Columbia, Manitoba, and Saskatchewan; Schedule II elsewhere).<sup>19,28</sup> <ul style="list-style-type: none"> <li>○ To find where to get naloxone in your province or territory go to <a href="https://www.canada.ca/en/health-canada/services/substance-use/problematic-prescription-drug-use/opioids/naloxone.html#5">https://www.canada.ca/en/health-canada/services/substance-use/problematic-prescription-drug-use/opioids/naloxone.html#5</a>.</li> </ul> </li> </ul>

Question	Answer/Pertinent Information/Suggested Resources
<b>What is the availability of naloxone nasal spray?</b>	<p><b>Naloxone nasal spray</b> 4 mg (the 2 mg strength, found in product labeling, is not marketed in the US or Canada). Generics are available in the US.</p> <ul style="list-style-type: none"><li>• <i>Narcan</i> 4 mg brand is available OTC (US only). <i>Narcan</i> (Canada) and generic 4 mg formulations (US only) are available without a prescription through various protocols (depending on the state/provincial/territorial laws).</li><li>• Each carton contains TWO blister-packed, single-dose nasal sprays, each containing 4 mg of naloxone.<sup>3,26</sup></li><li>• Be careful of mix-ups between the newer <i>Narcan</i> <b>nasal spray</b> and an older, no longer available <i>Narcan</i> brand <b>injection</b>. For some, “<i>Narcan</i>” has become synonymous with injectable naloxone.</li></ul> <p><b>Kloxxado nasal spray</b> (US only) is available as an 8 mg strength.<sup>25</sup></p> <ul style="list-style-type: none"><li>• Each carton contains TWO blister-packed, single-dose nasal sprays, each containing 8 mg of naloxone.<sup>25</sup></li></ul> <p>Naloxone kits for <b>intranasal administration using injectable solution</b>; each kit should contain:<sup>2,5</sup></p> <ul style="list-style-type: none"><li>• Two mucosal atomization devices (e.g., MAD Nasal device). Examples of sources for ordering include:<ul style="list-style-type: none"><li>○ Common Cents EMS Supplies: <a href="https://www.savelives.com/">https://www.savelives.com/</a> or 866-388-4599.</li><li>○ Teleflex Medical: <a href="https://www.teleflex.com/usa/en/index.html">https://www.teleflex.com/usa/en/index.html</a> (US) or <a href="https://www.teleflex.com/ca/en/index.html">https://www.teleflex.com/ca/en/index.html</a> (Canada) or 866-246-6990 (US) or 800-387-9699 (Canada).</li></ul></li><li>• Two naloxone 2 mg/2 mL Luer lock prefilled syringes (e.g., IMS/Amphastar [NDC# 76329-3369-1], Dr. Reddy’s [NDC# 43598-750-11], Auromedics [NDC# 55150-345-01]).<ul style="list-style-type: none"><li>○ Some atomizers contain needles allowing naloxone to be drawn up from vials instead of attaching to syringes.</li></ul></li></ul>
<b>How should commercially available naloxone or nalmefene nasal sprays be administered?</b>	<ul style="list-style-type: none"><li>• Keep the spray in the original packaging until use.<sup>3,25,26,34</sup></li><li>• The person to receive the spray should be lying on their back.<sup>3,25,26,34</sup></li><li>• Remove the spray from its packaging and hold it with the thumb on the bottom of the plunger and the first and middle fingers on either side of the nozzle.<sup>3,25,26,34</sup></li><li>• Tilt the patient’s head back gently. Insert the tip of the nozzle into one nostril until the fingers holding the nozzle are touching the bottom of the person’s nose.<sup>3,25,26,34</sup><ul style="list-style-type: none"><li>○ When administering naloxone in young children, if the nozzle does not fit completely in the nostril, make sure the nozzle seals the nostril before naloxone administration.<sup>26</sup></li></ul></li><li>• Press the plunger firmly, spraying the naloxone or nalmefene into the nostril.<sup>3,25,26,34</sup></li><li>• Remove the nozzle from the nostril.<sup>3,25,26,34</sup></li></ul>

Question	Answer/Pertinent Information/Suggested Resources
<b>How should an intranasal naloxone kit (using prefilled injectable naloxone) be administered?</b>	Follow these steps to <b>administer prefilled naloxone injectable solution intranasally</b> . <sup>5,30</sup> (There are three parts: the atomizer device, a plastic tube, and the naloxone Luer lock prefilled needleless syringes.) <ul style="list-style-type: none"><li>• Remove the two yellow caps from the plastic tube AND remove the cap from the naloxone.</li><li>• Hold the atomizer device by its plastic wings and attach it to the plastic tube by twisting it into place.</li><li>• Screw the naloxone Luer lock container into the barrel of the tube.</li><li>• Place atomizer into one of the patient’s nostrils.</li><li>• Deliver <b>the first half of the naloxone</b> dose (1 mL [1 mg]) by giving a short, vigorous push on the naloxone container.</li><li>• Move the atomizer into the patient’s other nostril.</li><li>• Deliver the <b>second half of the naloxone</b> dose (1 mL [1 mg]) by giving a short, vigorous push on the naloxone container.</li></ul>
<b>How should injectable naloxone be administered intramuscularly?</b>	Follow these steps to <b>administer naloxone IM</b> . <sup>4,16,30,32</sup> <ul style="list-style-type: none"><li>• Remove the cap from the naloxone vial or break the naloxone ampule neck.</li><li>• Uncover the needle.</li><li>• If using a naloxone ampule:<ul style="list-style-type: none"><li>○ Insert the needle into the liquid within the ampule.</li><li>○ Pull back the plunger to draw 1 mL (0.4 mg) into the syringe.</li></ul></li><li>• If using a naloxone vial:<ul style="list-style-type: none"><li>○ Hold the vial upside down and insert the needle through the rubber plug.</li><li>○ Pull back the plunger to draw 1 mL (0.4 mg) into the syringe.</li></ul></li><li>• Inject the naloxone into the muscle of the shoulder, thigh, or upper outer buttocks at a 90° angle.<ul style="list-style-type: none"><li>○ Feel comfortable giving the injection through clothing, if it is not thick, such as a jacket or heavy sweater.</li></ul></li><li>• Repeat doses can be given every two to three minutes if there is no change in the patient’s status or if the patient gets sleepy again.</li><li>• Naloxone needles used for IM injection should be disposed of in a sharps container.<ul style="list-style-type: none"><li>○ Emergency medical personnel might do this.<sup>16</sup></li></ul></li></ul>

Question	Answer/Pertinent Information/Suggested Resources
<b>How can injectable naloxone be used to prepare naloxone kits for administration via injection?</b>	<p><b>US:</b></p> <ul style="list-style-type: none"><li>• Naloxone for <b>IM injection</b> is available from single-dose or multidose naloxone vials. For <b>IM</b> use, it is recommended to give naloxone 0.4 mg (1 mL).<sup>4</sup> Each kit should contain:<sup>2,4</sup><ul style="list-style-type: none"><li>○ one multidose vial or two single-dose vials.</li><li>○ one 3 mL syringe with an appropriately sized needle (22 to 25 gauge, 1 to 1.5 inches). Include two syringes if using the single-dose vials and ten syringes with a 10 mL vial.</li></ul></li><li>• Information on preparing and prescribing naloxone rescue kits is available at:<ul style="list-style-type: none"><li>○ Harm Reduction Coalition: <a href="https://harmreduction.org/issues/overdose-prevention/naloxone-kits-materials/">https://harmreduction.org/issues/overdose-prevention/naloxone-kits-materials/</a>.</li><li>○ Prescribe to Prevent: <a href="http://prescribetoprevent.org/">http://prescribetoprevent.org/</a>.</li></ul></li></ul> <p><b>Canada:</b></p> <ul style="list-style-type: none"><li>• Injectable naloxone is available as 0.4 mg/mL in 1 mL single-dose ampoules or vials, and as 1 mg/mL in 2 mL multidose vials.<sup>16</sup></li><li>• The College of Pharmacists of British Columbia recommend dispensing at least two doses of 0.4 mg/mL naloxone plus at least two 3 mL <b>safety syringes</b> (e.g., <i>Vanish Point</i>, <i>BD Integra</i>) with 25 gauge, 1-inch needles.<sup>20</sup></li><li>• Other supplies that might be helpful include a one-way barrier breathing mask for giving rescue breaths and an ampoule breaker.<sup>20</sup><ul style="list-style-type: none"><li>○ Gloves and alcohol swabs could be included but are not necessary, as the injection can be given through lightweight clothing.<sup>16,20</sup></li></ul></li><li>• Kits are also available from <b>community-based programs</b> (often called “take-home naloxone” programs).<ul style="list-style-type: none"><li>○ These programs provide kits and naloxone training.<sup>21</sup></li><li>○ Kits typically consist of two 1 mL single-dose ampoules, needles, syringes, alcohol swabs, one-way barrier mask, instructions, and case.<sup>21</sup></li></ul></li><li>• For available programs throughout Canada, see:<ul style="list-style-type: none"><li>○ <a href="https://www.canada.ca/en/health-canada/services/opioids/naloxone.html#5">https://www.canada.ca/en/health-canada/services/opioids/naloxone.html#5</a>.</li><li>○ <a href="https://www.pharmacists.ca/cpha-ca/assets/File/cpha-on-the-issues/Naloxone_Scan_Nov_2021.pdf">https://www.pharmacists.ca/cpha-ca/assets/File/cpha-on-the-issues/Naloxone_Scan_Nov_2021.pdf</a>.</li></ul></li></ul>
<b>What is the maximum recommended naloxone dose?</b>	<ul style="list-style-type: none"><li>• There is not a well-established maximum naloxone dose.<sup>18</sup></li><li>• Product labeling indicates initial doses of 0.4 mg to 8 mg (dose based on route of administration), with repeat doses, as necessary.<sup>10,18</sup></li><li>• Typically, patients will respond to the first dose.<sup>18</sup><ul style="list-style-type: none"><li>○ Second doses are often supplied as a backup.<sup>18</sup></li><li>○ Additional doses may be needed, especially when emergency help is delayed, and initial naloxone doses wear off.<sup>18</sup></li><li>○ Higher doses or repeated administration may be required in patients having taken longer-acting opioids or partial opioid agonists (buprenorphine, pentazocine).<sup>11</sup></li></ul></li><li>• Healthy volunteers have received 24 mg without experiencing toxicity.<sup>18</sup></li></ul>

Question	Answer/Pertinent Information/Suggested Resources
<p><b>What happens after a dose of naloxone or nalmefene is administered?</b></p>	<ul style="list-style-type: none"> <li>• The duration of most opioids is longer than that of naloxone (30 to 120 minutes).<sup>6,23,24,30,32</sup> The duration of nalmefene (about six hours) is as long as most opioids.<sup>34</sup> <ul style="list-style-type: none"> <li>○ Emergency medical help should be requested (call 911) immediately.<sup>6,13,16,26,34</sup></li> </ul> </li> <li>• If symptoms return or if the patient doesn't respond or achieve the desired response (i.e., adequate spontaneous breathing), and emergency medical help has not yet arrived, repeat doses of naloxone can be given every two to three minutes (or nalmefene every two to five minutes).<sup>3,16,25,26,32,34</sup> <ul style="list-style-type: none"> <li>○ When giving additional doses of <i>Narcan</i> (Rx), <i>Kloxxado</i>, or <i>Opvee</i> nasal spray, use alternate nostrils.<sup>3,25,26,34</sup></li> </ul> </li> <li>• Rescue breathing may be required, and ideally, patients experiencing opioid overdose should be given oxygen.<sup>6,34</sup></li> <li>• Naloxone or nalmefene use may precipitate withdrawal in opioid-dependent patients. (See details above in section, "What should you tell patients, caregivers, and families about naloxone?")</li> <li>• If naloxone is given to a patient who is not opioid-dependent or is not opioid-intoxicated, it has no clinical effects.<sup>6</sup></li> </ul>
<p><b>How should naloxone and nalmefene be stored?</b></p>	<ul style="list-style-type: none"> <li>• Store naloxone and nalmefene at room temperature and protect from light.<sup>8,34</sup></li> <li>• <i>Kloxxado</i>, <i>Narcan</i>, and <i>Opvee</i> nasal sprays and <i>Zimhi</i> prefilled syringe should be stored in the original packaging. <ul style="list-style-type: none"> <li>○ <i>Narcan</i> nasal spray: Store below 77°F (25°C), but not below 5°F (-15°C), as it freezes and will not be usable if needed.<sup>3,26</sup> Excursions up to 104°F (40°C) are allowed.<sup>3,26</sup></li> <li>○ <i>Kloxxado</i> nasal spray: Store between 68°F and 77°F (20°C and 25°C). Excursions up to 104°F (40°C) and down to 41°F (5°C) are allowed. <i>Kloxxado</i> freezes below 5°F (-15°C) and will not spray; however, it can be thawed for 15 minutes at room temperature and then used. Seek emergency medical help right away if needed, and do NOT wait for <i>Kloxxado</i> to thaw.<sup>25</sup></li> <li>○ Store injectable naloxone between 15°C and 30°C (59°F and 86°F).<sup>16</sup></li> <li>○ <i>Zimhi</i> prefilled syringe: Store between 68°F and 77°F (20°C and 25°C).<sup>32</sup> Excursions between 59°F and 86°F (15°C and 30°C) are allowed.<sup>32</sup></li> <li>○ <i>Opvee</i> nasal spray: Store between 15°C and 25°C (59°F and 77°F). Short-term excursions between 4°C and 40°C (39°F and 104°F) are allowed.<sup>34</sup></li> </ul> </li> <li>• It is a good idea for patients to carry naloxone products with them, and to tell family and others who may need to administer naloxone where it is kept.<sup>3,25,32</sup> <ul style="list-style-type: none"> <li>○ While counseling patients about naloxone storage consider reminding them to keep their prescription opioid secure; divulging opioid use to others might invite theft.</li> </ul> </li> <li>• Patients should periodically check the appearance of their injectable naloxone.<sup>16,32</sup> <ul style="list-style-type: none"> <li>○ If the solution is discolored, cloudy, or contains particulates it should be replaced.<sup>16,32</sup></li> </ul> </li> <li>• Naloxone products (and syringes, if applicable) should be replaced <b>before</b> the expiration date.<sup>8</sup> <ul style="list-style-type: none"> <li>○ If stored properly, naloxone products should be effective at least until the manufacturer's expiration date. Typically, the shelf-life is 12 to 24 months.<sup>8,19</sup></li> <li>○ It has been suggested that pharmacists dispense naloxone with at least a six-month shelf-life at time of sale, and ideally longer than one year.<sup>20</sup></li> </ul> </li> </ul>

Question	Answer/Pertinent Information/Suggested Resources
<p><b>Can naloxone be provided to a third party?</b></p>	<ul style="list-style-type: none"> <li>• Providing naloxone or nalmefene to a third-party (e.g., to a caregiver or family member) as opposed to a patient may be permitted. <ul style="list-style-type: none"> <li>○ In the <b>US</b>, to find out about providing naloxone or nalmefene to a third-party, consult any of the following: <ul style="list-style-type: none"> <li>▪ <a href="http://pdaps.org/datasets/laws-regulating-administration-of-naloxone-1501695139">http://pdaps.org/datasets/laws-regulating-administration-of-naloxone-1501695139</a>.</li> <li>▪ your state medical board.</li> <li>▪ patient-specific insurance programs, including Medicaid and Medicare.</li> </ul> </li> <li>○ In <b>Canada</b>, naloxone is available to anyone without a prescription (unscheduled in Alberta, British Columbia, Manitoba, Saskatchewan; Schedule II elsewhere).<sup>19,28</sup></li> </ul> </li> </ul>
<p><b>How much do naloxone and nalmefene cost in the US?<sup>a</sup></b></p>	<ul style="list-style-type: none"> <li>• <b>OTC <i>Narcan</i> (naloxone) nasal spray 4 mg</b> (includes two nasal sprays): \$45.</li> <li>• <b>Naloxone nasal spray 4 mg</b> (includes one or two nasal sprays): ~\$102/box containing two sprays (generic). <ul style="list-style-type: none"> <li>○ For first responders, governments, schools, and community programs, <i>Narcan</i> may be available at a reduced cost (844-4NARCAN).</li> <li>○ Most major insurance plans cover naloxone nasal spray 4 mg.<sup>29</sup></li> </ul> </li> <li>• <b><i>Kloxxado</i> (naloxone) nasal spray 8 mg</b> (includes two nasal sprays): \$125.</li> <li>• <b><i>Obvee</i> (nalmefene) nasal spray 2.7 mg</b> (includes two nasal sprays): \$98</li>   <li>• <b>Naloxone prefilled injectable syringe 2 mg/2 mL</b>: \$31/2 mg syringe.</li> <li>• <b><i>Zimhi</i> (naloxone) prefilled injectable syringe 5 mg/0.5 mL</b>: \$62.50 for one or \$125 for two syringes.</li>   <li>• <b>Naloxone “kits” (US)</b>: <ul style="list-style-type: none"> <li>○ <b>Intranasal</b> (two naloxone 2 mg/2 mL prefilled syringes [~\$63] plus two nasal atomizers [~\$20]): ~\$84.</li> <li>○ <b>Intramuscular</b> (two naloxone 0.4 mg/mL single-dose vials): ~\$20 plus cost of syringes.</li> <li>○ Check with your local health department, as some may offer naloxone at no charge.</li> <li>○ Some insurance plans, including Medicaid and Medicare in some states, will cover the kits or some components.<sup>10</sup></li> </ul> </li> </ul>
<p><b>How much does naloxone cost in Canada?<sup>a</sup></b></p>	<ul style="list-style-type: none"> <li>• <b><i>Narcan</i> nasal spray</b>: \$157/box containing two sprays.</li> <li>• <b><i>S.O.S. Naloxone</i> (Canada)</b>: <ul style="list-style-type: none"> <li>○ naloxone 0.4 mg/mL vial or ampoule: ~\$15/0.4 mg dose.</li> <li>○ <i>BD Integra</i> syringe: ~\$0.80/syringe.</li> </ul> </li> <li>• Do not expect naloxone to be covered on the patient’s provincial/territorial drug plan, or their extended healthcare plan.</li> <li>• If cost is an issue, consider referring patients to a community-based program, which might provide naloxone for free. In some provinces, publicly funded take-home kits with injectable naloxone or <i>Narcan</i> nasal spray may be available through pharmacies. See <a href="https://www.pharmacists.ca/cpha-ca/assets/File/cpha-on-the-issues/Naloxone_Scan_Nov_2021.pdf">https://www.pharmacists.ca/cpha-ca/assets/File/cpha-on-the-issues/Naloxone_Scan_Nov_2021.pdf</a> or <a href="https://www.canada.ca/en/health-canada/services/opioids/naloxone.html#5">https://www.canada.ca/en/health-canada/services/opioids/naloxone.html#5</a>.</li> </ul>



Question	Answer/Pertinent Information/Suggested Resources
<p><b>How do I bill for naloxone- or nalmefene-related counseling (US)?</b></p>	<p><b>Prescribers (US)</b></p> <ul style="list-style-type: none"> <li>• Use the codes for Screening, Brief Intervention, and Referral to Treatment (SBIRT) to bill for counseling a patient about how to recognize overdose and how to administer naloxone or nalmefene.<sup>6</sup> <ul style="list-style-type: none"> <li>○ Use these billing codes for SBIRT:<sup>6</sup> <ul style="list-style-type: none"> <li>▪ CPT 99408 (commercial insurance, 15 to 30 minutes).</li> <li>▪ CPT 99409 (commercial insurance, longer than 30 minutes).</li> <li>▪ G0396 (Medicare, 15 to 30 minutes).</li> <li>▪ G0397 (Medicare, longer than 30 minutes).</li> <li>▪ H0049 (Medicaid, alcohol and/or drug screening).</li> <li>▪ H0050 (Medicaid, alcohol and/or drug services, brief intervention, per 15 minutes).</li> </ul> </li> <li>○ For counseling and instruction on the safe use of opioids, including the use of naloxone or nalmefene, outside of the context of SBIRT services, the prescriber should document the time spent and use the E&amp;M code that accurately captures the time and complexity. For example, in new patients deemed appropriate for opioid pharmacotherapy, when a substantial and appropriate amount of additional time is used to provide a separate service such as behavioral counseling (e.g., opioid overdose risk assessment and naloxone or nalmefene administration training), consider using modifier –25 in addition to the E&amp;M code.<sup>6</sup></li> </ul> </li> <li>• When using an evidence-based opioid misuse/abuse screening tool, CPT Code 99420 (administration and interpretation of health risk assessment instrument) can be used for patients with commercial insurance.<sup>6</sup></li> </ul>
<p><b>Does the availability of opioid receptor blockers encourage opioid misuse?</b></p>	<ul style="list-style-type: none"> <li>• In communities where naloxone distribution programs exist, opioid overdose deaths decrease.<sup>14</sup> These programs have not been shown to increase drug use but do increase interest in treatment.<sup>6</sup></li> <li>• Patients who are given a naloxone prescription have positive behavioral changes related to opioids, such as improved dosing (e.g., being more careful to take the right dose, and count the hours between doses), and improved knowledge of opioids and overdose.<sup>12</sup></li> </ul>
<p><b>Are there liability issues related to opioid receptor blockers?</b></p>	<ul style="list-style-type: none"> <li>• The medico-legal risks of prescribing naloxone to opioid users are low.<sup>6,10</sup></li> <li>• In the US, laws are being drafted and passed to protect prescribers, dispensers, and bystanders who administer naloxone. <ul style="list-style-type: none"> <li>○ See <a href="http://pdaps.org/datasets/laws-regulating-administration-of-naloxone-1501695139">http://pdaps.org/datasets/laws-regulating-administration-of-naloxone-1501695139</a> for information by state.</li> <li>○ Good Samaritan Laws may require “good faith” and “reasonable care.”</li> </ul> </li> <li>• In Canada, the “Good Samaritan Drug Overdose Act” became law in May of 2017.<sup>22</sup> <ul style="list-style-type: none"> <li>○ Witnesses are protected and encouraged to carry and provide naloxone during suspected opioid overdoses.<sup>22</sup></li> </ul> </li> </ul>

Question	Answer/Pertinent Information/Suggested Resources
<b>How are nalmefene (US only) and naloxone used in hospitals and emergency departments?</b>	<ul style="list-style-type: none"><li>• IM and subcutaneous <b>naloxone</b> are the most likely methods of naloxone administration in hospitals and emergency departments.<ul style="list-style-type: none"><li>○ Subcutaneous and IM doses have a similar onset of action and are less expensive than intranasal naloxone.<sup>18</sup></li><li>○ The onset of action for subcutaneous and IM naloxone is likely quicker than obtaining IV access.<sup>18</sup></li></ul></li><li>• <b>Nalmefene</b> (US only) injection is for administration by healthcare professionals, with a similar cost<sup>a</sup> (\$30/2 mg vial) compared to naloxone.<sup>7,24,31</sup> There is no evidence that it is more effective than naloxone, and its longer duration of action may be a downside (e.g., longer duration of stay for monitoring and management of precipitated withdrawal).<ul style="list-style-type: none"><li>○ Nalmefene can be given IV (fastest onset), IM, or subcutaneously.<sup>7</sup> The IV dose is 0.5 mg/70 kg, with a second dose of 1 mg/70 kg two to five minutes later if needed.<sup>7</sup> If precipitated withdrawal is a concern, consider a test dose of 0.1 mg/70 kg, followed in two minutes by the usual dose if there are no withdrawal symptoms.<sup>7</sup></li></ul></li><li>• Follow your facility policies for providing naloxone or nalmefene at discharge. Options may include:<ul style="list-style-type: none"><li>○ Providing naloxone, naloxone kits, or nalmefene directly to patients, families, or caregivers.</li><li>○ Giving a written order for patients, families, or caregivers to obtain naloxone or nalmefene (e.g., pharmacy, health department).</li></ul></li></ul>

**Abbreviations:** CNS = central nervous system; IM = intramuscular; IV = intravenous; OTC = over-the-counter.

a. Pricing based on wholesale acquisition cost (WAC). US medication pricing by Elsevier, accessed November 2023.

Users of this resource are cautioned to use their own professional judgment and consult any other necessary or appropriate sources prior to making clinical judgments based on the content of this document. Our editors have researched the information with input from experts, government agencies, and national organizations. Information and internet links in this article were current as of the date of publication.

### Levels of Evidence

In accordance with our goal of providing Evidence-Based information, we are citing the **LEVEL OF EVIDENCE** for the clinical recommendations we publish.

Level	Definition	Study Quality
<b>A</b>	Good-quality patient-oriented evidence.*	<ol style="list-style-type: none"> <li>High-quality randomized controlled trial (RCT)</li> <li>Systematic review (SR)/Meta-analysis of RCTs with consistent findings</li> <li>All-or-none study</li> </ol>
<b>B</b>	Inconsistent or limited-quality patient-oriented evidence.*	<ol style="list-style-type: none"> <li>Lower-quality RCT</li> <li>SR/Meta-analysis with low-quality clinical trials or of studies with inconsistent findings</li> <li>Cohort study</li> <li>Case control study</li> </ol>
<b>C</b>	Consensus; usual practice; expert opinion; disease-oriented evidence (e.g., physiologic or surrogate endpoints); case series for studies of diagnosis, treatment, prevention, or screening.	

\***Outcomes that matter to patients** (e.g., morbidity, mortality, symptom improvement, quality of life).

[Adapted from Ebell MH, Siwek J, Weiss BD, et al. Strength of recommendation taxonomy (SORT): a patient-centered approach to grading evidence in the medical literature. *Am Fam Physician*. 2004 Feb 1;69(3):548-56.

<https://www.aafp.org/pubs/afp/issues/2004/0201/p548.html>.]

### References

- FDA news release. FDA requiring labeling changes for opioid pain medicines, opioid use disorder medicines regarding naloxone. July 23, 2020. <https://www.fda.gov/news-events/press-announcements/fda-requiring-labeling-changes-opioid-pain-medicines-opioid-use-disorder-medicines-regarding>. (Accessed November 6, 2023).
- National Harm Reduction Coalition. How to assemble naloxone kits. Last modified April 22, 2022. <https://harmreduction.org/issues/overdose-prevention/naloxone-kits-materials/>. (Accessed November 6, 2023).
- Product information Narcan nasal spray. Emergent Devices. Plymouth Meeting, PA 19462. November 2020.
- Prescribe to Prevent. Naloxone for overdose prevention (intramuscular). [http://www.prescribetoprevent.org/wp-content/uploads/2012/11/one-pager\\_22.pdf](http://www.prescribetoprevent.org/wp-content/uploads/2012/11/one-pager_22.pdf). (Accessed November 6, 2023).
- Prescribe to Prevent. Naloxone for overdose prevention (intranasal). <http://www.prescribetoprevent.org/wp-content/uploads/2012/11/naloxone-one-pager-in-nov-2012.pdf>. (Accessed November 6, 2023).
- Substance Abuse and Mental Health Services Administration. SAMHSA opioid overdose prevention toolkit. HHS publication no (SMA) 18-4742. Rockville, MD: Substance Abuse and Mental Health Services Administration, June 2018. <https://store.samhsa.gov/product/Opioid-Overdose-Prevention-Toolkit/SMA18-4742>. (Accessed November 6, 2023).
- Prescribing information for nalmefene. Purdue Pharma. Stamford, CT 06901. February 2022.
- College of Psychiatric & Neurologic Pharmacists. Naloxone access: a practical guideline for pharmacists. Updated February 2015. <http://prescribetoprevent.org/wp2015/wp-content/uploads/naloxone-access.pdf>. (Accessed November 6, 2023).
- Prescribe to Prevent. Instruction for healthcare professionals: prescribing naloxone. [http://www.prescribetoprevent.org/wp-content/uploads/2012/11/one-pager\\_12.pdf](http://www.prescribetoprevent.org/wp-content/uploads/2012/11/one-pager_12.pdf). (Accessed November 6, 2023).
- Prescribe to Prevent. FAQ. <https://prescribetoprevent.org/faq-2/>. (Accessed November 6, 2023).
- Jordan MR, Morrisonponce D. Naloxone. [Updated April 29, 2023]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2023 Jan-.
- Behar E, Rowe C, Santos GM, et al. Primary Care Patient Experience with Naloxone Prescription. *Ann Fam Med*. 2016 Sep;14(5):431-6.
- CDC. Lifesaving naloxone. April 21, 2023. <https://www.cdc.gov/stopoverdose/naloxone/index.html>. (Accessed November 6, 2023).
- Wheeler E, Jones TS, Gilbert MK, et al. Opioid Overdose Prevention Programs Providing Naloxone to Laypersons - United States, 2014. *MMWR Morb Mortal Wkly Rep*. 2015 Jun 19;64(23):631-5.
- Green TC, Davis C, Xuan Z, et al. Laws Mandating Coprescription of Naloxone and Their Impact on Naloxone Prescription in Five US States, 2014-2018. *Am J Public Health*. 2020 Jun;110(6):881-887.
- Product monograph for S.O.S. Naloxone Hydrochloride Injection. Sandoz Canada. Boucherville, QC JB 7K8. April 2017.
- College of Pharmacists of British Columbia. Checklist for naloxone training. February 28, 2023. [http://library.bcpharmacists.org/6\\_Resources/6-5\\_Pharmacy\\_Resources/5184-](http://library.bcpharmacists.org/6_Resources/6-5_Pharmacy_Resources/5184-)

- Naloxone\_Checklist\_For\_Training.pdf. (Accessed November 6, 2023).
18. Pammatt R. Naloxone for opioid overdose – virtual Q&A. Canadian Pharmacy Association: pharmacy practice webinar. [https://www.pharmacists.ca/cpha-ca/assets/File/education-practice-resources/NaloxoneWebinarQA\\_Sept22.pdf](https://www.pharmacists.ca/cpha-ca/assets/File/education-practice-resources/NaloxoneWebinarQA_Sept22.pdf). (Accessed November 6, 2023).
  19. Canadian Agency for Drugs and Technologies in Health. Funding and management of naloxone programs in Canada. January 26, 2023. <https://www.cadth.ca/funding-and-management-naloxone-programs-canada-0#public-drug-plans-coverage-for-naloxone>. (Accessed November 6, 2023).
  20. College of Pharmacists of British Columbia. Community pharmacy distribution of naloxone. [http://library.bcpharmacists.org/6\\_Resources/6-5\\_Pharmacy\\_Resources/5187-Naloxone\\_Training.pdf](http://library.bcpharmacists.org/6_Resources/6-5_Pharmacy_Resources/5187-Naloxone_Training.pdf). (Accessed November 6, 2023).
  21. Canadian Centre on Substance Abuse. The availability of take-home naloxone in Canada. CCENDU Bulletin. March 2016. <https://ccsa.ca/sites/default/files/2019-05/CCSA-CCENDU-Take-Home-Naloxone-Canada-2016-en.pdf>. (Accessed November 6, 2023).
  22. Government of Canada. About the Good Samaritan Drug Overdose Act. Date modified November 23, 2021. <https://www.canada.ca/en/health-canada/services/substance-use/problematic-prescription-drug-use/opioids/about-good-samaritan-drug-overdose-act.html>. (Accessed November 6, 2023).
  23. National Institute on Drug Abuse. Naloxone drug facts. January 2022. <https://nida.nih.gov/publications/drugfacts/naloxone>. (Accessed November 6, 2023).
  24. Edinoff AN, Nix CA, Reed TD, et al. Pharmacologic and clinical considerations of nalmefene, a long duration opioid antagonist, in opioid overdose. *Psychiatry Int* 2021;365-78.
  25. Product information for Kloxxado. Hikma Specialty USA. Columbus, OH 43228. May 2022.
  26. Product monograph for Narcan nasal spray. Emergent BioSolutions Canada. Oakville, ON L6L 0C4. March 2021.
  27. American Medical Association. AMA opioid task force. Help save lives: co-prescribe naloxone to patients at risk of overdose. August 2017. <https://www.end-oid-epidemic.org/wp-content/uploads/2017/08/AMA-Opioid-Task-Force-naloxone-one-pager-updated-August-2017-FINAL.pdf>. (Accessed November 6, 2023).
  28. Canadian Pharmacists Association. Publicly-funded take-home naloxone in pharmacies across Canada. November 2021. [https://www.pharmacists.ca/cpha-ca/assets/File/cpha-on-the-issues/Naloxone\\_Scan\\_Nov\\_2021.pdf](https://www.pharmacists.ca/cpha-ca/assets/File/cpha-on-the-issues/Naloxone_Scan_Nov_2021.pdf). (Accessed November 6, 2023).
  29. Narcan. Frequently asked questions. <https://www.narcan.com/frequently-asked-questions/>. (Accessed April 4, 2023).
  30. National Harm Reduction Coalition. Opioid overdose basics: responding to opioid overdose. Last modified September 1, 2020. <https://harmreduction.org/issues/overdose-prevention/overview/overdose-basics/responding-to-opioid-overdose/>. (Accessed November 6, 2023).
  31. Purdue Pharma. Purdue Pharma introduces nalmefene HCL injection, 2 mg/2mL (1 mg/mL) in the U.S. for treatment of known or suspected overdose with natural or synthetic opioids. June 21, 2022, <https://www.purduepharma.com/news/2022/06/21/purdue-pharma-introduces-nalmefene-hcl-injection-2mg-2ml-1mg-1ml-in-the-u-s-for-the-treatment-of-known-or-suspected-overdose-with-natural-or-synthetic-opioids/>. (Accessed November 6, 2023).
  32. Product information for Zimhi. USWM. Louisville, KY 40241. October 2021.
  33. FDA. FDA approves first over-the-counter naloxone nasal spray. March 29, 2023. <https://www.fda.gov/news-events/press-announcements/fda-approves-first-over-counter-naloxone-nasal-spray>. (Accessed November 6, 2023).
  34. Product information for Opvee. Indivior. North Chesterfield, VA 23235. June 2023.
  35. State of Oregon. Joint statement on distribution of short-acting opioid antagonists (naloxone and nalmefene). 2023. [https://www.oregon.gov/osbn/Documents/IS\\_JointStatement\\_Naloxone.pdf](https://www.oregon.gov/osbn/Documents/IS_JointStatement_Naloxone.pdf). (Accessed November 6, 2023).
  36. Alabama Public Health. Standing order of the state health officer opioid reversal agent distribution for overdose prevention. 2023. <https://www.alabamapublichealth.gov/pharmacy/assets/standingorder.pdf>. (Accessed November 6, 2023).

***Cite this document as follows: Clinical Resource, Meds for Opioid Overdose. Pharmacist's Letter/Pharmacy Technician's Letter/Prescriber Insights. November 2023. [391108]***

—To access hundreds more clinical resources like this one, visit [trchealthcare.com](https://trchealthcare.com) to log in or subscribe—